

Online Technical Appendix for Fenton, Wyper, McCartney, Minton, 'Socioeconomic inequality in recent mortality trends in Scotland'

Jon Minton

Introduction

This document describes all data, methods and code used to generate the results within the above-listed article.

All data are publically available, and you are free to interrogate and use the code in your own analyses.

All data preparation, analyses, figures and tables were generated using the R programming language, with the exception of some initial data preparation done within Microsoft Excel. This data preparation only involved changing the structure of existing data within an Excel sheet, rather than the contents of such data.

Data

The data used are from [this National Records of Scotland \(NRS\) webpage](#). The dataset 'Latest tables based on 2013 ESP' (European Standardized Population) was downloaded in Excel format.

All analyses presented in the paper were based on table 7 of this Excel Workbook.

The contents of table 7 of the Workbook were rearranged so as to conform with data structure recommended by (Wickham, (2014) 'Tidy Data')(<https://www.jstatsoft.org/article/view/v059i10/v59i10.pdf>), producing a new datasheet, called `flat_data`, which was imported directly into R. All further processing and analyses were conducted within R.

Workflow and results

The following code chunks perform the analyses which are presented in the paper.

Where additional information are required as to the methods, they are described adjacent to the code chunks.

Loading the required packages

```
pacman::p_load(tidyverse, readxl, cowplot, kableExtra)
```

Note: `pacman` is an R package for managing R packages. It has to be installed once using `install.packages("pacman")`, but once installed will either install or load other packages as required.

Load data

```
dta <- read_excel("data/ASMR_SIMD_2001_2017_indexed_trends.xlsx", sheet = "flat_data")
```

Tidying the data

```
names(dta) <- c("year", "overall", "q1", "q2", "q3", "q4", "q5", "gender")

dta_tidy <- dta %>%
  gather(key = "simd", value = "asmr", -year, -gender) %>%
  mutate(SIMD = factor(simd,
    levels = c("q1", "q2", "q3", "q4", "q5", "overall"),
    labels = c("Q.1 (Most deprived)",
               "Q.2", "Q.3", "Q.4", "Q.5 (Least deprived)", "Overall")
  )
)
```

Visualising

This is one part of the first figure. The R package `ggplot2` is used to produce this visualisation.


```

p1 <- dta_tidy %>%
  ggplot(aes(x = year )) +

  facet_wrap(~gender) +
  geom_line(aes(y = asmr, group = SIMD, linetype = SIMD, size = SIMD, color = SIMD)) +
  scale_size_manual(values = c(1, 1.2, 1, 1.2, 1, 1.4)) +
  scale_linetype_manual(values = c(2,3,4,5, 6, 1)) +
  scale_color_manual(values = c("black", "grey", "black", "grey", "black", "blue")) +
  labs(x = "Year", y = "Age Standardised mortality rate per 100 000") +
  scale_y_continuous(expand = c(0, 0), limits = c(0, 2500), minor_breaks = seq(0, 2500, by = 100)) +
  scale_x_continuous(minor_breaks = 2001:2017) +
  geom_vline(xintercept = 2012, linetype = "dashed") +
  geom_vline(xintercept = 2006, linetype = "dashed") +
  annotate("text", y = 100, x = 2006 + (2012 - 2006) / 2, label = "2006 to 2011") +
  annotate("text", y = 100, x = 2012 + (2018 - 2012) / 2, label = "2012 to 2017") +
  geom_ribbon(
    aes(x = year, ymin = q5, ymax =q1),
    alpha = 0.1, fill ="red",
    data = dta_tidy %>% filter(simd %in% c("q1", "q5")) %>% select(-SIMD) %>% spread(simd, asmr)
  ) +
  background_grid(major = "xy", minor = "xy")

```

p1



The following chunk calculates the percentage change, from the earlier to latter period, in ASMR by gender and SIMD quintile.

```

percent_changes <- dta_tidy %>%
  mutate(period = case_when(
    between(year, 2012, 2017) ~ "2012-2017",
    between(year, 2006, 2011) ~ "2006-2011",
    TRUE ~ NA_character_) %>% factor(levels = c("2012-2017", "2006-2011"))) %>%
  group_by(gender, simd, period) %>%
  filter(year == min(year) | year == max(year)) %>%
  filter(!is.na(period)) %>%
  group_by(gender, SIMD, period) %>%
  summarise(percent_change = - 100 * (1 - asmr[year == max(year)] / asmr[year == min(year)])) %>%
  ungroup()

```

```

## Warning: Factor `period` contains implicit NA, consider using
## `forcats::fct_explicit_na`

```


Note: The contents of `percent_changes` look as follows:

```
percent_changes
```

Note: The code chunk above can be readily adapted to showing absolute change (for example), as follows

```
dta_tidy %>%
  mutate(period = case_when(
    between(year, 2012, 2017) ~ "2012-2017",
    between(year, 2006, 2011) ~ "2006-2011",
    TRUE ~ NA_character_) %>% factor(levels = c("2012-2017", "2006-2011"))) %>%
  group_by(gender, simd, period) %>%
  filter(year == min(year) | year == max(year)) %>%
  filter(!is.na(period)) %>%
  group_by(gender, SIMD, period) %>%
  summarise(
    percent_change = - 100 * (1 - asmr[year == max(year)] / asmr[year == min(year)]),
    absolute_change = asmr[year==max(year)] - asmr[year == min(year)] # additional line
  ) %>%
  ungroup() %>% # indicative visualisation
  ggplot(aes(x = SIMD, y = absolute_change)) +
  geom_point() +
  facet_grid(period~gender) +
  geom_hline(yintercept = 0) +
  theme(axis.text.x = element_text(angle = 90))
```

```
## Warning: Factor `period` contains implicit NA, consider using
## `forcats::fct_explicit_na`
```




The following shows the second part of the figure in the paper

```

p2 <- percent_changes %>%
  mutate(SIMD = factor(SIMD,
    levels = c("Q.1 (Most deprived)",
               "Q.2", "Q.3", "Q.4", "Q.5 (Least deprived)", "Overall"),
    labels = c("Most", "Q2", "Q3", "Q4", "Least", "Overall")
  )
) %>%
filter(SIMD != "Overall") %>%
ggplot(aes(x = SIMD, y = percent_change, group = period, shape = period)) +
facet_wrap(~ gender) +
geom_point(size = 5) +
stat_smooth(method = "lm", se = F) + # This produces the blue line with the regression slopes
geom_hline(yintercept = 0) +
geom_hline( # This adds the overall percent change
  aes(yintercept = percent_change, group = period),
  data = percent_changes %>% filter(SIMD == "Overall"),
  linetype = "dashed"
) +
theme(axis.text.x = element_text(angle = 90)) +
labs(y = "Percent change in ASMR by period", x = "SIMD Quintile") +
scale_shape_manual("Period", values = c(17, 16))

```

p2



Combined figure


The following produces the combined figure comprising the two parts shown previously. The figure is rendered as a png format image at 300dpi, and placed in the directory 'figures'.

```

p_both <- plot_grid(p1, p2, labels = c("A", "B"), ncol = 1, align = "v")

p_both

```



```
ggsave("figures/combined_figure_2012.png", dpi = 300, units = "cm", height = 30, width = 30)
```

Table

The following shows how the contents of the table were produced.

These summarise a series of univariate linear regressions of SIMD quintile against percentage change in ASMR within distinct periods.

The columns `intercept` and `gradient` present summary information about point estimates and 95% confidence intervals for the intercept and gradient of the the regressions, which are then presented in the formatted table.

A functional programming approach was adopted in order to produce the same analyses consistently for different gender and period combinations, using functions within the (`purrr` package)[<https://purrr.tidyverse.org/>].

```

get_ci <- function(x){
  tmp <- x %>% summary() %>% coefficients()

  return(
    list(
      lower = tmp[,1] - 1.96 * tmp[,2],
      upper = tmp[,1] + 1.96 * tmp[,2]
    )
  )
}

# Model parameters
tbl_1 <- percent_changes %>%
  filter(SIMD != "Overall") %>%
  mutate(qnt = unclass(SIMD) - 1) %>% # This is so the intercept refers to the 1st quintile (not the 'zeroth'
  ' quintile)
  select(gender, period, percent_change, qnt) %>%
  group_by(gender, period) %>%
  nest() %>%
  mutate(mdl = map(data, ~lm(percent_change ~ qnt, data = .x))) %>%
  mutate(`R. sq.` = map_dbl(mdl, ~summary(.x)[`r.squared`][1])) %>%
  mutate(gradient = map_dbl(mdl, ~coef(.x)[`qnt`])) %>%
  mutate(intercept = map_dbl(mdl, ~coef(.x)[`(Intercept)`])) %>%
  mutate(cis = map(mdl, get_ci)) %>%
  mutate(
    int_lower = map_dbl(cis, ~.[`lower`][1]),
    int_upper = map_dbl(cis, ~.[`upper`][1]),
    grd_lower = map_dbl(cis, ~.[`lower`][2]),
    grd_upper = map_dbl(cis, ~.[`upper`][2])
  ) %>%
  select(gender, period, `R. sq.`,
    gradient, grd_lower, grd_upper,
    intercept, int_lower, int_upper
  ) %>%
  mutate(
    gradient = paste0(
      format(round(gradient, 2), nsmall = 2),
      " (",
      format(round(grd_lower, 2), nsmall = 2),
      ", ",
      format(round(grd_upper, 2), nsmall = 2),
      ") "
    )
  ) %>%
  mutate(
    intercept = paste0(
      format(round(intercept, 2), nsmall = 2),
      " (",
      format(round(int_lower, 2), nsmall = 2),
      ", ",
      format(round(int_upper, 2), nsmall = 2),
      ") "
    )
  ) %>%
  select(-grd_lower, -grd_upper, -int_lower, -int_upper)

tbl_1

tbl_2 <- percent_changes %>% spread(SIMD, percent_change)

tbl_both <- inner_join(tbl_2, tbl_1)

## Joining, by = c("gender", "period")

tbl_both

```

The following converts the above table into a more neatly formatted table using the `kable` and `kableExtra` packages.

```
tbl_both %>%
  mutate(period = factor(period, levels = c("2006-2011", "2012-2017"))) %>%
  arrange(gender, period) %>%
  knitr::kable(
    digits = 2,
    caption = "Percent change in ASMR by gender, SIMD quintile, and period"
  ) %>%
  kableExtra::kable_styling() %>%
  kableExtra::add_header_above(c(" ", " ", "Percentages" = 6, "Model results" = 3)) %>%
  kableExtra::footnote("Overall: Whole of Scotland. R.Sq. : R-Squared for model. Gradient: Increase in % cha
nge per unit increase in quintile. Intercept: Predicted % change in most deprived quintile. For gradient and
intercept, values in parentheses show lower and upper 95% confidence intervals of coefficients respectively.
")
```

Percent change in ASMR by gender, SIMD quintile, and period

| gender | period | Percentages | | | | | | Model results | | |
|--------|-----------|---------------------|-------|-------|-------|----------------------|---------|---------------|----------------------|------------------------|
| | | Q.1 (Most deprived) | Q.2 | Q.3 | Q.4 | Q.5 (Least deprived) | Overall | R. sq. | gradient | intercept |
| Female | 2006-2011 | -7.21 | -7.37 | - | - | -7.81 | -10.11 | 0.12 | -0.70 (-2.84, 1.44) | -8.38 (-13.62, -3.14) |
| Female | 2012-2017 | 0.69 | -2.65 | -2.96 | -5.53 | -7.79 | -3.46 | 0.96 | -1.98 (-2.44, -1.53) | 0.32 (-0.80, 1.44) |
| Male | 2006-2011 | -9.81 | - | - | - | -12.57 | -11.82 | 0.79 | -0.63 (-1.00, -0.26) | -10.19 (-11.09, -9.29) |
| Male | 2012-2017 | 2.11 | -2.27 | -2.78 | -3.57 | -4.44 | -2.00 | 0.80 | -1.44 (-2.25, -0.63) | 0.69 (-1.29, 2.67) |
| Total | 2006-2011 | -8.56 | -9.18 | - | - | -9.04 | -10.59 | 0.12 | -0.40 (-1.60, 0.81) | -9.44 (-12.39, -6.48) |
| Total | 2012-2017 | 1.35 | -2.32 | -2.38 | -4.61 | -6.05 | -2.59 | 0.93 | -1.71 (-2.25, -1.17) | 0.62 (-0.70, 1.93) |

Note:
Overall: Whole of Scotland. R.Sq. : R-Squared for model. Gradient: Increase in % change per unit increase in quintile. Intercept: Predicted % change in most deprived quintile. For gradient and intercept, values in parentheses show lower and upper 95% confidence intervals of coefficients respectively.

Approach discussed in sensitivity analysis

The sensitivity analysis to the paper showed the effect of using the fitted values for the first and last year in each of the periods, rather than the values themselves. This approach can address any concern that the first and last year within either period were in any way anomalous or uncharacteristic of change within the period as a whole.

The function `broom::augment` was used to extract fitted values for each year within each period, gender and SIMD combination. The fitted values, `.fitted`, were then used in place of the observed values, `asmr`, as in the main analyses.

```
percent_changes_pred <- dta_tidy %>%
  mutate(period = case_when(
    between(year, 2012, 2017) ~ "2012-2017",
    between(year, 2006, 2011) ~ "2006-2011",
    TRUE ~ NA_character_) %>% factor(levels = c("2012-2017", "2006-2011"))) %>%
  group_by(gender, SIMD, simd, period) %>%
  nest() %>%
  mutate(mdl = map(data, ~lm(asmr ~ year, data = .x))) %>%
  mutate(aug = map(mdl, broom::augment)) %>%
  select(-data, -mdl) %>%
  unnest() %>%
  filter(!is.na(period)) %>%
  group_by(gender, SIMD, period) %>%
  filter(year == min(year) | year == max(year)) %>%
  summarise(percent_change = - 100 * (1 - .fitted[year == max(year)] / .fitted[year == min(year)])) %>%
  ungroup()
```

```
## Warning: Factor `period` contains implicit NA, consider using
## `forcats::fct_explicit_na`

## Warning: Factor `period` contains implicit NA, consider using
## `forcats::fct_explicit_na`
```

```
percent_changes_pred_overall <- percent_changes_pred %>%
  group_by(gender, period) %>%
  summarise(percent_change = mean(percent_change)) %>%
  ungroup() %>%
  mutate(SIMD = "Mean of quintiles") %>%
  select(gender, SIMD, period, percent_change)

percent_changes_pred <- percent_changes_pred %>%
  bind_rows(percent_changes_pred_overall) %>%
  mutate(SIMD = factor(SIMD,
    levels = c("Q.1 (Most deprived)",
               "Q.2", "Q.3", "Q.4", "Q.5 (Least deprived)", "Overall", "Mean of quintile
s"),
    labels = c("Most", "Q2", "Q3", "Q4", "Least", "Overall", "Mean of quintiles")
  )
)
```

```
## Warning in bind_rows(x, .id): binding factor and character vector,
## coercing into character vector
```


```
## Warning in bind_rows(x, .id): binding character and factor vector,
## coercing into character vector
```

The subfigure using this alternative stratgy is therefore produced as follows:

```
p2a <- percent_changes_pred %>%
  filter(!(SIMD %in% c("Overall", "Mean of quintiles"))) %>%
  ggplot(aes(x = SIMD, y = percent_change, group = period, shape = period)) +
  facet_wrap(~ gender) +
  geom_point(size = 5) +
  stat_smooth(method = "lm", se = F) +
  geom_hline(yintercept = 0) +
  geom_hline(
    aes(yintercept = percent_change, group = period),
    data = percent_changes_pred %>% filter(SIMD == "Overall"),
    linetype = "dashed"
  ) +

  theme(axis.text.x = element_text(angle = 90)) +
  labs(y = "Percent change in ASMR by period", x = "SIMD Quintile") +
  scale_shape_manual("Period", values = c(17, 16))

p2a
```

```
ggsave("figures/figla_using_alt_method_2012.png", dpi = 300, units = "cm", height = 16, width = 30)
```

The table of the above, using this alternative modelling strategy, is produced using the code chunks below

```

get_ci <- function(x){
  tmp <- x %>% summary() %>% coefficients()

  return(
    list(
      lower = tmp[,1] - 1.96 * tmp[,2],
      upper = tmp[,1] + 1.96 * tmp[,2]
    )
  )
}

# Model parameters
tbl_1a <- percent_changes_pred %>%
  filter(!(SIMD %in% c("Overall", "Mean of quintiles"))) %>%
  mutate(qnt = unclass(SIMD) - 1) %>% # This is so the intercept refers to the 1st quintile (not the 'zeroth'
  'quintile)
  select(gender, period, percent_change, qnt) %>%
  group_by(gender, period) %>%
  nest() %>%
  mutate(mdl = map(data, ~lm(percent_change ~ qnt, data = .x))) %>%
  mutate(`R. sq.` = map_dbl(mdl, ~summary(.x)[`r.squared`][1])) %>%
  mutate(gradient = map_dbl(mdl, ~coef(.x)[`qnt`])) %>%
  mutate(intercept = map_dbl(mdl, ~coef(.x)[`Intercept`])) %>%
  mutate(cis = map(mdl, get_ci)) %>%
  mutate(
    int_lower = map_dbl(cis, ~.[`lower`][1]),
    int_upper = map_dbl(cis, ~.[`upper`][1]),
    grd_lower = map_dbl(cis, ~.[`lower`][2]),
    grd_upper = map_dbl(cis, ~.[`upper`][2])
  ) %>%
  select(gender, period, `R. sq.`,
    gradient, grd_lower, grd_upper,
    intercept, int_lower, int_upper
  ) %>%
  mutate(
    gradient = paste0(
      format(round(gradient, 2), nsmall = 2),
      " (",
      format(round(grd_lower, 2), nsmall = 2),
      ", ",
      format(round(grd_upper, 2), nsmall = 2),
      ") "
    )
  ) %>%
  mutate(
    intercept = paste0(
      format(round(intercept, 2), nsmall = 2),
      " (",
      format(round(int_lower, 2), nsmall = 2),
      ", ",
      format(round(int_upper, 2), nsmall = 2),
      ") "
    )
  ) %>%
  select(-grd_lower, -grd_upper, -int_lower, -int_upper)

tbl_1a

```

```

tbl_2a <- percent_changes_pred %>% filter(SIMD != "Mean of quintiles") %>% spread(SIMD, percent_change)

tbl_both_a <- inner_join(tbl_2a, tbl_1a)

```

```
## Joining, by = c("gender", "period")
```

```
tbl_both_a
```

And the table using the alternative approach

```
tbl_both_a %>%
  mutate(period = factor(period, levels = c("2006-2011", "2012-2017"))) %>%
  arrange(gender, period) %>%
  knitr::kable(
    digits = 2,
    caption = "Percent change in ASMR by gender, SIMD quintile, and period. (Alternative method)"
  ) %>%
  kableExtra::kable_styling() %>%
  kableExtra::add_header_above(c(" ", " ", "Percentages" = 6, "Model results" = 3)) %>%
  kableExtra::footnote("Overall: Whole of Scotland. R.Sq. : R-Squared for model. Gradient: Increase in % cha
nge per unit increase in quintile. Intercept: Predicted % change in most deprived quintile. For gradient and
intercept, values in parentheses show lower and upper 95% confidence intervals of coefficients respectively.
")
```

Percent change in ASMR by gender, SIMD quintile, and period. (Alternative method)

| gender | period | Percentages | | | | | | Model results | | |
|--------|-----------|-------------|--------|--------|--------|--------|---------|---------------|----------------------|-------------------------|
| | | Most | Q2 | Q3 | Q4 | Least | Overall | R. sq. | gradient | intercept |
| Female | 2006-2011 | -7.35 | -9.09 | -13.32 | -13.59 | -9.76 | -10.94 | 0.29 | -0.93 (-2.58, 0.72) | -8.76 (-12.80, -4.72) |
| Female | 2012-2017 | 1.60 | -2.38 | -0.75 | -4.50 | -7.60 | -2.47 | 0.85 | -2.05 (-3.03, -1.07) | 1.38 (-1.03, 3.78) |
| Male | 2006-2011 | -11.90 | -13.14 | -12.62 | -13.06 | -12.40 | -13.11 | 0.08 | -0.09 (-0.44, 0.26) | -12.44 (-13.29, -11.59) |
| Male | 2012-2017 | 2.69 | -1.92 | -0.42 | -5.06 | -3.76 | -1.40 | 0.71 | -1.60 (-2.78, -0.43) | 1.51 (-1.36, 4.39) |
| Total | 2006-2011 | -9.63 | -10.73 | -12.82 | -12.76 | -10.45 | -11.67 | 0.16 | -0.37 (-1.31, 0.58) | -10.55 (-12.85, -8.24) |
| Total | 2012-2017 | 2.22 | -1.85 | -0.19 | -4.54 | -5.73 | -1.72 | 0.83 | -1.86 (-2.80, -0.92) | 1.70 (-0.60, 4.01) |

Note:
Overall: Whole of Scotland. R.Sq. : R-Squared for model. Gradient: Increase in % change per unit increase in quintile. Intercept: Predicted % change in most deprived quintile. For gradient and intercept, values in parentheses show lower and upper 95% confidence intervals of coefficients respectively.